

GROUPS OF INTERMEDIATE GROWTH

LAURENT BARTHOLDI

ABSTRACT. I describe a class of groups acting on rooted trees that all have intermediate word growth between polynomial and exponential.

The argument constructs a functional equation on the growth formal power series, and derives the growth properties from its analytical behaviour.

This places under a unified proof all known or conjectured examples of such groups, and answers a few open questions by Rostislav Grigorchuk.

I then sketch some algebraic results enjoyed by this class of groups, and list what seem promising open questions and further directions.

1. INTRODUCTION

Let G be a group, finitely generated by a set S . The *word norm* on G is the norm

$$(1) \quad \|g\| = \min \{n \in \mathbb{N} \mid g = s_1 \dots s_n \text{ for some } s_i \in S\}.$$

The *growth function* of G is the function $\gamma(n) = \#\{g \in G \mid \|g\| \leq n\}$, measuring the growth of the volume of balls in the metric space G .

Say two functions γ, δ are equivalent if $\gamma(n) \leq \delta(Cn)$ and $\delta(n) \leq \gamma(Cn)$ for some large enough $C \in \mathbb{N}$. Then although γ depends both on G and S , its equivalence class depends only on G .

This notion of growth was introduced in the 1950's in the USSR [Šva55] and in the 1960's in the West [Mil68a], in relation with properties of Riemannian manifolds and coverings : assume G is the fundamental group of a compact metrized CW -complex K . Then the growth function of G measures the growth of balls in the universal cover $\tilde{K}$.

If γ is dominated by a polynomial function, i.e. $\gamma(n) \leq Cn^D$ for some $C, D \in \mathbb{N}$, then G is of *polynomial growth*. This happens when G is free abelian (indeed then $\tilde{K}$ is Euclidean space), and more generally when G has a nilpotent subgroup of finite index, as was observed by John Milnor [Mil68a] and Joseph Wolf [Wol68]. Mikhail Gromov proved conversely that these are the only examples [Gro81].

In the other end of the spectrum, if $\gamma(n) \geq C^n$ for some $C > 1$, then G is of *exponential growth*. Clearly, one must have $C \leq \#S$. This happens e.g. when G is non-virtually-nilpotent linear [Tit72] or non-elementary hyperbolic [GH90].

John Milnor asked in 1968 [Mil68b] whether there exist finitely generated groups of *intermediate growth*, i.e. with growth greater than polynomial but less than exponential. The first example of such a group was discovered in 1978 by Rostislav

Date: February 1, 2008.

1991 *Mathematics Subject Classification.* **20E08** (Groups acting on trees), **20F32** (Geometric group theory).

Key words and phrases. intermediate growth; growth of groups; tree automorphisms.

The author acknowledges support from the "Swiss National Fund for Scientific Research", stipend #83R-063208.

Grigorchuk [Gri83, Gri91]. He later constructed a continuum of groups of intermediate growth [Gri84], and showed that the growth functions γ of all these groups satisfy

$$e^{\sqrt{n}} \lesssim \gamma(n) \not\sim e^n;$$

he also showed that the growth function of his main example, the “first Grigorchuk group” [Har00, Chapter VIII], satisfies

$$\gamma(n) \lesssim e^{n^\beta}, \quad \beta = \log_{32}(31) \approx 0.991.$$

I have improved the upper [Bar98] and lower [Bar01] bounds of this group to

$$e^{n^{0.5157}} \lesssim \gamma(n) \lesssim e^{n^{0.7675}};$$

and have obtained with Zoran Šuník [BŠ01] upper and lower bounds for the continuum of Grigorchuk groups.

The Grigorchuk groups are, under a very small condition, infinite torsion groups. A similar construction of infinite torsion groups was proposed by Narain Gupta and Said Sidki [GS83a, GS83b], and the natural question as to whether these groups are also of intermediate growth has been long open. I address it in this paper by the following result:

Data. Let $\Sigma = \{1, \dots, d\}$ be a finite set; the set Σ^* of finite sequences over Σ is naturally (the vertex set of) a d -regular rooted tree, rooted at the empty sequence, if one connects for all $\sigma_i \in \Sigma$ the vertices $\sigma_1 \dots \sigma_n$ and $\sigma_1 \dots \sigma_n \sigma_{n+1}$.

Let A be a subgroup of the symmetric group of Σ ; let B be a finite group, and let $\tilde{B} = B_1 \times B_2 \times \dots$ be the direct product of countably many copies of B ; let $\phi_1, \dots, \phi_{d-1}$ be homomorphisms $\tilde{B} \rightarrow A$, and let $\phi_d : \tilde{B} \rightarrow \tilde{B}$ be the one-sided shift $(b_1, b_2, \dots) \mapsto (1, b_1, b_2, \dots)$. Write ϕ_i^n for the restriction of ϕ_i to B_n .

Action. A and $\tilde{B}$ act on Σ^* as follows: the action of $a \in A$ is

$$(\sigma_1 \sigma_2 \dots \sigma_n)^a = (\sigma_1)^a \sigma_2 \dots \sigma_n,$$

and the action of $\tilde{b} \in \tilde{B}$ is defined inductively by $()^{\tilde{b}} = ()$ and

$$(\sigma_1 \sigma_2 \dots \sigma_n)^{\tilde{b}} = \sigma_1 (\sigma_2 \dots \sigma_n)^{\phi_{\sigma_1}(\tilde{b})}.$$

Let G be the subgroup of $\text{Aut } \Sigma^*$ generated by A and B_1 , i.e. the largest quotient of $A * B_1$ that acts faithfully on Σ^* . Such a group G is called a *splitter-mixer group*, since its action is two-step: the $\tilde{b}$ ’s “split” in actions of a ’s, and the actions of a ’s “mix” the subtrees below them.

Theorem 1.1. *Let G be a splitter-mixer group as defined above. Then*

1. G has subexponential growth;
2. assume that A acts transitively on Σ ; that for all $1 \neq b \in B$ there exists infinitely many $(n, i) \in \mathbb{N} \times \{1, \dots, d-1\}$ with $\phi_i^n(b) \neq 1$; and that $\langle \phi_1^n(B), \dots, \phi_{d-1}^n(B) \rangle = A$ for all $n \in \mathbb{N}$. Then G has superpolynomial growth, unless $A = 1$ or $A = B = \mathbb{Z}/2$.

The description of groups of intermediate growth followed a long evolution. They were first described as transformations of the interval $[0, 1]$, and only later as permutations of a rooted tree, finally presented algebraically as Σ^* .

Theorem 1.1 proves that the following groups have intermediate growth:

The first Grigorchuk group: It is obtained by taking $\Sigma = \{1, 2\}$; $A = \{1, a = (1, 2)\}$; $B = A \times A = \{1, b, c, d\}$; and

$$\phi_1^n(b) = \phi_1^{n-1}(c) = \phi_1^{n-2}(d) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{3}, \\ a & \text{else.} \end{cases}$$

The “Grigorchuk Overgroup”: It is obtained by taking $\Sigma = \{1, 2\}$; $A = \{1, a = (1, 2)\}$; $B = A^3 = \langle \tilde{b}, \tilde{c}, \tilde{d} \rangle$; and

$$\phi_1^n(\tilde{b}) = \phi_1^{n-1}(\tilde{c}) = \phi_1^{n-2}(\tilde{d}) = \begin{cases} a & \text{if } n \equiv 1 \pmod{3}, \\ 1 & \text{else.} \end{cases}$$

Note that this group $\tilde{G}$ contains the first Grigorchuk group as the subgroup $\langle a, \tilde{b}\tilde{c}, \tilde{b}\tilde{d}, \tilde{c}\tilde{d} \rangle$. The growth of $\tilde{G}$ is shown to be exponential in [BG02].

The Grigorchuk groups G_ω : These groups form an uncountable family described in [Gri84], indexed by $\omega \in \{X, Y, Z\}^{\mathbb{N}}$. They are obtained by taking $\Sigma = \{1, 2\}$; $A = \{1, a\}$; $B = A \times A$; and $\phi_1^n = \omega_n$, if one interprets X, Y, Z as the three non-trivial homomorphisms $B \rightarrow A$.

Grigorchuk proves in [Gri84] that if every letter appears infinitely often in ω , then G_ω is an infinite torsion group of intermediate growth.

The Gupta-Sidki group: It is obtained by taking $\Sigma = \{1, 2, 3\}$, $A = \langle x = (1, 2, 3) \rangle$; $B = A$; and $\phi_1^n(x) = \phi_2^n(x^{-1}) = x$.

The Gupta-Sidki p -groups: They are defined in [GS83a], and in our setting are obtained by taking $\Sigma = \{1, 2, \dots, p\}$ for $p \geq 3$, $A = \langle x = (1, 2, \dots, p) \rangle$, $B = A$, and $\phi_1^n(x) = \phi_2^n(x^{-1}) = x$, $\phi_3^n(x) = \dots = \phi_{p-1}^n(x) = 1$. These groups are infinite p -torsion groups for all p .

The Gupta-Fabrykowski group: It is obtained by taking $\Sigma = \{1, 2, 3\}$, $A = \langle x = (1, 2, 3) \rangle$; $B = A$; and $\phi_1^n(x) = \phi_2^n(x) = x$. This group was announced to be of intermediate growth in [FG85], but the arguments presented failed to convince, and were improved in [FG91].

The “Spinal groups”: These groups were first introduced in [GS84] where they were given with a condition for torsion. They were rediscovered in [BŠ01] with a condition for word and period growth. In the present setting, they amount to imposing $\phi_2^n = \dots = \phi_{d-1}^n(b) = 1$, and the condition that for all $b \in B$, we have $\phi_1^n(b) = 1$ for infinitely many n . Note that in compensation for these restrictions some quantitative growth estimates are obtained.

The “Square group”: It is an example proposed by Grigorchuk, who stated its growth as an interesting open question. One takes $\Sigma = \{1, 2, 3, 4\}$, $A = \langle x = (1, 2, 3, 4) \rangle$; $B = A$; and $\phi_1^n(x) = x$, $\phi_2^n(x) = 1$, $\phi_3^n(x) = x^{-1}$.

The paper is organized as follows: Section 2 recalls notions on growth of groups. Section 3 and 4 prove respectively the first and second part of Theorem 1.1. Section 5 describes the distribution of the projections for some examples of splitter-mixer groups, Section 6 gives some algebraic properties of splitter-mixer groups (torsion, branchness), and Section 7 closes with some remarks and open questions.

The emphasis of this paper is on growth; a future paper will study more deeply the algebraic nature of the groups introduced here.

This work was initiated when I was a PhD student of Professor Pierre de la Harpe in the University in Geneva. I thank Pierre heartily for everything he has taught me during those years.

2. GROWTH

Let G be a group, finitely generated as a monoid by a set S . The word norm (1) gives rise to the growth function $\gamma(n) = \#\{g \in G \mid \|g\| \leq n\}$, conveniently studied through its *growth series*, the generating function

$$\Gamma(z) = \sum_{g \in G} z^{\|g\|} = (1 - z) \sum_{n \geq 0} \gamma(n) z^n.$$

This power series has been extensively studied, in particular when viewed as an analytic function in the ball of radius $1/\#S$. If G is virtually abelian [Ben83], hyperbolic [GH90, Chapter 9], or a Coxeter group [Par93], then Γ is a rational function of z . If G is the Heisenberg group $G = \langle x, y \mid [x, y] \text{ central} \rangle$, its growth series is also rational [Sha89]. There are other nilpotent groups whose growth series is rational for some generating sets and irrational for others [Sto96]; the same phenomenon occurs for other cyclic fiber groups [Sha94].

Note that if G is finite, then Γ is a polynomial; otherwise, the convergence radius of Γ is at least $1/(\#S - 1)$, and at most 1. The following is a formulation of the root test:

Lemma 2.1. *G has subexponential growth if and only if Γ converges in the unit disk.*

The following result is due to Pierre Fatou and Fritz Carlson:

Lemma 2.2 ([Fat06], page 368; [Car21]). *If Γ is a formal power series with integer coefficients, converging in the unit disk, then Γ is either rational or transcendental; in the latter case, it has the unit disk as natural boundary.*

Corollary 2.3. *If G has intermediate growth, then Γ is a transcendental function.*

Proof. By Lemmata 2.2 and 2.1, Γ is either rational or transcendental. If Γ were rational, it would have finitely many singularities on the unit circle, and each of them would be of pole type $f(z) = (z - \omega)^{-D}$. Then G would have polynomial growth, of degree $D + 1$ for the largest such D . $\square$

Lemma 2.4. *Let a_n, b_n, c_n be real sequences related by the equation*

$$c_n = \sum_{i+j=n} \binom{n}{i} a_i b_j.$$

Let $\Gamma_a, \Gamma_b, \Gamma_c$ be the related generating series $\Gamma_a = \sum_{n \geq 0} a_n z^n$, etc. Then for all $\epsilon > 0$ we have

$$\Gamma_c(z) = \frac{1}{2i\pi} \oint_{|w|=\epsilon} \Gamma_a((1+w)z) \Gamma_b((1+w^{-1})z) \frac{dw}{w}.$$

In consequence, if Γ_a and Γ_b respectively have radius of convergence $1/\alpha$ and $1/\beta$, then Γ_c has radius of convergence at least $1/(\alpha + \beta)$.

Proof. By linearity, it suffices to check the formula for $\Gamma_a(z) = z^i$ and $\Gamma_b(z) = z^j$. Then

$$\Gamma_c(z) = \frac{z^{i+j}}{2i\pi} \oint_{|w|=\epsilon} (1+w)^i (1+w^{-1})^j \frac{dw}{w} = \binom{i+j}{j} z^{i+j},$$

by the Cauchy residue formula.

Next, consider the integral on the circle $|w| = \epsilon = \beta/\alpha$. If $|z| < 1/(\alpha + \beta)$, then $|(1+w)z| < 1/\alpha$ and $|(1+w^{-1})z| < 1/\beta$, so the integral converges uniformly. $\square$

Corollary 2.5. *Let $\eta > 0$ be fixed, and assume the real sequences a_n, b_n, c_n are related by the equation¹*

$$c_n = \sum_{i+j=n} \eta \binom{\eta n}{\eta i} a_i b_j.$$

Let $\Gamma_a, \Gamma_b, \Gamma_c$ be the related generating series $\Gamma_a = \sum_{n \geq 0} a_n z^n$, etc., with radius of convergence $1/\alpha, 1/\beta, 1/\gamma$. Then

$$\gamma \geq \left(\alpha^{-1/\eta} + \beta^{-1/\eta} \right)^{-\eta}.$$

Proof. Consider the associated series $\Delta_a = \sum_{n \geq 0} a_n z^{\lfloor \eta n \rfloor}$, etc.; then the radius of convergence of Δ_a is $\alpha^{1/\eta}$, etc., and the power series $\Delta_a, \Delta_b, \Delta_c$ satisfy the hypotheses of the previous Lemma. $\square$

3. SUBEXPONENTIAL GROWTH

Let $A, \tilde{B} = B_1 \times \dots$ and ϕ_i^k be as in Theorem 1.1. For $k \in \mathbb{N}$, define G_k as the subgroup of $\text{Aut } \Sigma^*$ generated by A and B_k . In particular, $G = G_1$, and we will show that all G_k 's have subexponential growth.

As generating set S_k of G_k , we choose all symbols $s_{a,b,a'}$ with $a, a' \in A$ and $b \in B_k$. There is an obvious cancellation among these generators:

$$s_{a,b,a'} s_{a'',b',a'''} = s_{a,bb',a'''} \text{ if } a' a'' = 1,$$

and that is the only one that we shall use.

Given $w \in S_{k-1}^*$, we may uniquely write w as an expression $[w_1, \dots, w_d]a$, for some $a \in A$ and $w_1, \dots, w_d \in S_k^*$, as follows:

- first rewrite $w = s_{a'_1, b_1, a''_1} \dots s_{a'_n, b_n, a''_n}$ as a word $w = b_1^{a_1} \dots b_n^{a_n} a$, by setting $a_i = a'_1 a''_1 \dots a'_i$ for all i ;
- then rewrite each $b_i^{a_i}$ as $[\phi_{1^{a_i}}^{k-1}(b_i), \dots, \phi_{d^{a_i}}^{k-1}(b_i)]$, and multiply coordinate-wise;
- finally multiply out all consecutive elements of A , and rewrite each coordinate $w_i = a_1 b_1 \dots a_n b_n a_{n+1}$ as $s_{a_1, b_1, 1} \dots s_{a_n, b_n, a_{n+1}}$.

We insist that cancellations of the form $s_{a,b,a'} s_{(a')^{-1}, b^{-1}, a''} = aa''$ are never to be performed; the maximal cancellation is $s_{a,1,a''}$. We call the expression $[w_1, \dots, w_d]a$ the *decomposition* of w .

Our choice of generators has the following interpretation: the *length* of an element $g \in G_k$ is the minimal number of B -generators needed to express g . The total length of the w_i is the length of w minus the number of cancellations performed.

Fix an integer n , and consider a random word of length n . We say it has *uniform distribution* if all its B -symbols are uniform, independent variables in B , and all its A -symbols are non-trivial.

Lemma 3.1. *Let w be uniform of length n , and let $[w_1, \dots, w_d]a$ be its decomposition. Then each w_i is uniform.*

¹The binomial coefficients are evaluated on non-integers by the formula $\binom{n}{m} = \Gamma(n+1)/\Gamma(m+1)\Gamma(n-m)$.

Proof. Each w_i is a sequence of B -letters separated by strings of A -letters, and each of these letters is the ϕ -image of a uniformly distributed B -letter.

The A -letters are combined together by cancellation; but thanks to our assumption that trivial B -letters are not cancelled, the different strings of A -letters (which need not be uniform) never combine, so remain separated by products of uniform B -letters, which are again uniform. $\square$

We now wish to estimate the cancellation that occurs in the decomposition of w . In the strings of A -letters comprising w_i , each is the ϕ -image of a uniform B -letter. It follows that each such A -letter has probability at least $1/\#A$ of being trivial, in which case a cancellation occurs; unless it is the first or last letter, which does not contribute to the length of w_i . Therefore if w_i had length n_i before cancellation, then at worst² the probability that there be m cancellations in w_i is

$$(2) \quad C_m^{n_i-1} = \binom{n_i-1}{m} \left(\frac{1}{\#A} \right)^m \left(\frac{\#A-1}{\#A} \right)^{n_i-1-m}.$$

We now estimate the total length of the w_i . It is at most $|w| + d - 1 - m$, where m is the total number of cancellations. We may not claim that the cancellations on all w_i are independent — indeed, it may happen that abundant cancellation in w_i implies abundant cancellation in some other w_j .

However, the longest of the w_i has cancellation given by the previous paragraph, and the other w_j 's have cancellation correlated (to some positive or negative degree) with w_i . Call $p(m)$ the probability that there be m cancellations. If there is no correlation, then at worst $p(m) = C_m^{|w|-d+1}$, since then all $n-d+1$ inner A -syllables have independent probability at least $1/\#A$ of being trivial. If there is full positive correlation, then at worst $p(m) = \frac{1}{d} C_{m/d}^{|w|/d-1}$, since the d words $w_1, \dots, w_d$ can be assumed to behave in the same way.

We may therefore assume that there is an unknown parameter η , depending on G , such that at worst $p(m) = \eta C_{\eta m}^{\eta(|w|-d+1)}$.

The decomposition map on words induces a natural map $G_{k-1} \rightarrow G_k \wr_{\Sigma} A$, also written $g \mapsto [g_1, \dots, g_d]g_a$, where g_a is obtained by restricting the action of $g \in G_{k-1}$ to Σ , and g_i are obtained by restricting the action of gg_a^{-1} to $i\Sigma^*$. For all $k \in \mathbb{N}$ let Γ_k denote the growth series of G_k , and write $\langle n, \Gamma \rangle$ for the degree- n coefficient of Γ .

²In this section, “at worst” means “giving the largest growth estimate”.

We relate Γ_{k-1} to Γ_k as follows:

$$\begin{aligned}
\Gamma_{k-1}(z) &= \sum_{n \geq 0} \langle n, \Gamma_{k-1} \rangle z^n = \sum_{g \in G_{k-1}} z^{\|g\|} \\
&\leq \sum_{[g_1, \dots, g_d] a \in G_k \wr_\Sigma A} z^{\|[g_1, \dots, g_d] a\|} \\
&\leq \#A \sum_{[g_1, \dots, g_d] \in G_k \times \dots \times G_k} z^{\|[g_1, \dots, g_d]\|} \\
&\leq \#A \sum_{\substack{[g_1, \dots, g_d] \in G_k \times \dots \times G_k \\ 0 \leq m \leq \|[g_1, \dots, g_d]\|}} \eta C_m^\eta (\|[g_1, \dots, g_d]\| - d + 1) z^{\|[g_1, \dots, g_d]\|} \\
&\leq \#A \sum_{\substack{n_1, \dots, n_d, m \geq 0 \\ n + d - 1 - m = n_1 + \dots + n_d}} \eta C_m^{\eta(n-d+1)} \langle n_1, \Gamma_k \rangle \dots \langle n_d, \Gamma_k \rangle z^n \text{ with } n_i = \|g_i\| \\
(*) \quad &= z^d \#A \sum_{m, p \geq 0} \eta \binom{\eta(m+p)}{\eta m} \left(\frac{z}{\#A^\eta} \right)^m \left\langle p, z^{-2} \Gamma_k \left(z \left(\frac{\#A-1}{\#A} \right)^\eta \right)^d \right\rangle z^p.
\end{aligned}$$

Let $1/\rho_k$ be the convergence radius of Γ_k . Define the power series $\Delta(z) = (1 - z/\#A^\eta)^{-1}$, with radius of convergence $\#A^\eta$. Corollary 2.5 may be applied to the inequality (*) involving the power series Γ_k and Δ ; hence we obtain

$$\rho_{k-1} \leq \rho_k \frac{\#A-1}{\#A} + \frac{1}{\#A},$$

and hence by induction

$$\rho_k \leq (\rho_\ell - 1) \left(\frac{\#A-1}{\#A} \right)^{k-\ell} + 1 \text{ for all } k, \ell \in \mathbb{N}.$$

Since $\rho_\ell \leq \#S - 1$ for all $\ell \in \mathbb{N}$, we obtain $\rho_k = 1$ for all $k \in \mathbb{N}$. Therefore by Lemma 2.1 all G_k have subexponential growth.

For simplicity assume now $\eta = 1$. We actually derived more information from the previous computation; namely, writing $\Delta(z) = 1/(1 - z/\#A)$,

$$(3) \quad \Gamma_{k-1}(z) \leq \#A \frac{1}{2i\pi} \oint_{|w|=\epsilon} \Delta((1+w^{-1})z) \Gamma_k \left((1+w)z \frac{\#A-1}{\#A} \right)^d \frac{dw}{w}.$$

4. SUPERPOLYNOMIAL GROWTH

Make the assumptions of the second part of Theorem 1.1, and consider also the groups G_k introduced in the beginning of Section 3. The hypotheses show that the map $G_{k-1} \rightarrow G_k \wr_\Sigma A$ is surjective onto A and each of the factors G_k .

Assume for contradiction that some G_k has polynomial growth, say of degree D , and assume D is minimal among the growth rates of all G_n 's.

Consider the subgroup H of G_k , that fixes Σ pointwise. Restriction of the action to $d\Sigma^*$ defines a surjective map $H \rightarrow G_{k+1}$. On the other hand, consider the two cases:

- $d \geq 3$:** Then among the conjugates b^a for all $b \in B_k$ and $a \in A \setminus \text{Stab}_A(d)$, there must be at least two distinct, say $b_1^{a_1}$ and $b_2^{a_2}$, with same d -coordinate. Consider K , the normal closure in H of $b_1^{a_1} b_2^{-a_2}$.

d = 2: Then $A = \langle(1, 2)\rangle$ and $|B| > |A|$ by assumption, so there must be a non-trivial $b \in B_k$ with $\phi_1(b) = 1$. Consider K , the normal closure in H of $b^{(1,2)}$.

Since A acts transitively on Σ and all B_n 's map onto A , the group G_k acts transitively on Σ^n for all $n \in \mathbb{N}$. The $b^h \neq 1$ for $b \in B_k$ and $h \in H$ have an “axis”, which is the infinite path $(dd \dots)^h$ in Σ^* . Since H is transitive on $i\Sigma^*$ for all $i \in \Sigma$, it follows that there are infinitely many conjugates b^h , and hence in all cases K is infinite.

Clearly the restriction of K to $d\Sigma^*$ is trivial, so we have a surjective map $H \rightarrow G_{k+1}$ with infinite kernel. The growth of G_{k+1} is at least polynomial of degree D , so the growth of H , and hence of G_k , is at least polynomial of degree $D + 1$. This contradicts our assumption on the growth of G_k .

5. THE DISTRIBUTION OF THE DECOMPOSITION

The proof of subexponential growth made two simplifying assumptions: Given a uniform word w and its decomposition $[w_1, \dots, w_d]a$, each A -syllable of w_i has probability $1/\#A$ of being trivial, and these probabilities are independent, even between syllables in w_i and w_j coming from the same B -letter of w .

These simplifications were legitimate in that they only increased the growth estimate. We consider in this section whether they actually hold.

Proposition 5.1. *Assume w is a uniform word with decomposition $[w_1, \dots, w_d]a$. If all homomorphisms ϕ_i^n are onto, then the w_i are mutually independent uniform words.*

Proof. Clearly all the B -letters in $\{w_1, \dots, w_d\}$ are uniform and independent, since they come from all different B -letters in w .


Consider an arbitrary A -syllable in a word w_i . Since the ϕ_i^n are onto, it is a product of at least one uniform variable on A , and hence it is uniformly distributed.

Consider next two A -syllables in w_i and w_j . If $i = j$, then they are independent by Lemma 3.1. If $i \neq j$, then the B -letters from which they come in w are not all the same. Consider a B -letter contributing to the first syllable but not the next; then the distribution of the first syllable is a convolution with a uniform distribution over A that is independent of the second syllable, so these syllables have independent distribution. $\square$

The first group of intermediate growth for which the proposition does not apply is the “Fabrykowski-Gupta group”. There I found experimentally that on average the total length of the w_i 's is $\frac{1}{2}|w|$, and not $\frac{2}{3}|w|$.

For a splitter-mixer group G , consider a uniform word w of length n , and decompose it as $[w_1, \dots, w_d]a$. Consider the distribution of $|w_1| + \dots + |w_d|$. For n large, it will be a sum of many independent events, and hence will follow a normal distribution, with mean $\mu_G n$ and variance $\sigma_G n$.

The model used in section 3 made the assumptions $\mu_G = (\#A - 1)/\#A$ and $\sigma_G = \mu_G(1 - \mu_G)$. Figure 5 shows the distributions for the first examples of splitter-mixer groups, given by $A = B = \mathbb{Z}/d$ and $A = (\mathbb{Z}/2)^2, B = \mathbb{Z}/2$. They first are identified by a sequence of integers $\epsilon_1, \dots, \epsilon_{d-1}$ with $0 \leq \epsilon_i < d$, and are constructed as follows: $A = B = \mathbb{Z}/d\mathbb{Z} = \langle a \rangle$, and $\phi_i^n(a) = a^{\epsilon_i}$. The same construction gives the second, where $(\mathbb{Z}/2)^2$ is described as $\{1, i, j, k\}$ with $1 = i^2 = j^2 = k^2 = ijk$.



If there were no correlation between the coordinates, the distribution would follow the binomial law (2), with mean $\frac{\#A-1}{\#A}$ and variance $\mu_G(1 - \mu_G)$. The correlation factor η satisfies $\sigma_G\eta = \mu_G(1 - \mu_G)$; hence points above the parabola indicate groups with positive correlation, while points below the parabola indicate groups with negative correlation between the coordinates w_i .

The data in Figure 5 were obtained either by theoretical means (for G , $\tilde{G}$ and $d \leq 3$), or by Monte-Carlo simulation with 500 000 words of length 10 000 (for the 18 groups with $d = 4$ listed below).

Not all sequences of ϵ_i need be considered: the two sequences $\epsilon_1, \dots, \epsilon_{d-1}$ and $\epsilon_{d-1}, \dots, \epsilon_1$ define the same group, and so does $d-1-\epsilon_1, \dots, d-1-\epsilon_d$. These transformations amount to replacing the generator of A or B by its inverse.

We also restrict to sequences $\epsilon_1, \dots, \epsilon_{d-1}$ such that $\text{l.c.m.}(\epsilon_i) = 1$; this amounts to asking the embedding $G_{k-1} \rightarrow G_k \wr A$ to be subdirect.

The first examples are:

- 1 for $d = 2$; this is the infinite dihedral group;
- 0, 1 and 1, 1 and 1, 2 for $d = 3$; these are respectively the Fabrykowski-Gupta group, the Gupta-Sidki group, and the Bartholdi-Grigorchuk group. These 3 groups are non-isomorphic: the G-S group is torsion, the F-G is virtually torsion-free branch, and the B-G group is not branch.
- for $d = 4$, there are at most 18 isomorphism classes of groups, namely

001, 010, 021, 012, 102, 011, 013, 122, 212, 101, 103, 112, 132, 121, 123, 111, 113, 131.

I do not know which are isomorphic, but possibly their values of μ and σ serve as invariants to distinguish them.

- for $B = \mathbb{Z}/2$ and $A = (\mathbb{Z}/2)^2$, there is an action of $SL_2(\mathbb{Z}/2) = S_6$ on the sequences that simultaneously permutes the three entries and their contents. There are at most 9 isomorphism classes, namely

$$1ij, 1ik, 1jk, \quad iij, iik, jii, \quad ijk, jik, jki.$$

Note finally that the contraction distribution in a mixer-splitter group may be even larger than described here. First, the A -letters of the restrictions w_i need not be uniformly distributed over $A \setminus \{1\}$, even if those of w were. Secondly, there may be some extra cancellation coming from consideration of the B -letters.

6. MORE ON SPLITTER-MIXER GROUPS

In this section I give some more algebraic results on mixer-splitter groups. They extend naturally considerations by Taras Vovkivsky [Vov00] and Gupta and Sidki [GS84].

As above, we assume a mixer-splitter group G has been fixed, with data $\Sigma = \{1, \dots, d\}$; a group A acting on Σ ; and B and $\phi_i^n : B \rightarrow A$ for all $i \in \{1, \dots, d-1\}$ and $n \in \mathbb{N}$.

Let us make the assumption that the $\phi_i^n : B \rightarrow A$ are periodic functions, i.e. there is $k_i \in \mathbb{N}$ with $\phi_i^{n+k_i} = \phi_i^n$. Up to replacing Σ by a cartesian power of itself and extending A and B appropriately, we may suppose in that case that $\phi_i^{n+1} = \phi_i^n$ for all n, i . We will call G *regular* in that case.

Construct the graph $\mathfrak{G}$ with vertex set $B \times A$; for all $(b, a) \in V(\mathfrak{G})$, let $\{d, x_2, \dots, x_e\}$ be the Σ -orbit of a containing d , and call $e(b, a)$ its size. Put a directed edge from (a, b) to $\pi(b, a) = (b', a')$, where the action on $d\Sigma^*$ of $(ba)^e$ is $b'a'$. (Note that $b' = b$ or $b' = 1$, and $e(b, a) = 1$ implies $e\pi(b, a) = 1$).

Theorem 6.1. *Let G be a regular mixer-splitter group with A abelian. Then G is a torsion group if and only if all the cycles in $\mathfrak{G}$ are loops at vertices (b, a) with $e(b, a) = 1$.*

Proof. First, assume that the path starting at (b, a) does not reach a loop with $e(b, a) = 1$. Consider the elements $g_0 = ba$ and $g_{i+1} = g_i^{e^{\pi^i}(b, a)}$. Then g_i fixes the vertex d^i but not d^{i+1} , so all these elements are distinct and hence G is not torsion.

On the other hand, suppose that the only loops in $\mathfrak{G}$ are at (b, a) with $e(b, a) = 1$. Let $g = b_1 a_1 \dots b_k a_k$ be an element of G ; we reason by induction on k .

If $k = 1$, we have $g = ba$; let $\{d = x_1, x_2, \dots, x_e\}$ be the a -orbit of d in Σ . Assume first $e(b, a) = 1$; then the order of g is the l.c.m. of the order of b and the order of a , and hence is finite.

If $e(b, a) > 1$, then $g^{e(b, a)}$ fixes all subtrees $x_i \Sigma^*$, and acts like a conjugate of $b'a'$ on them. Since A is finite, a power of g^e furthermore fixes Σ and acts trivially on all subtrees not of this form. Repeating from (b', a') , we reach the previous case after a finite number of steps.

Suppose now $k > 1$, and assume all words of length at most $k - 1$ are torsion. Write the decomposition of g as $[g_1, \dots, g_d]a$, and let $\{d = x_1, x_2, \dots, x_e\}$ be the a -orbit of d in Σ . Then g^e fixes the subtree $d\Sigma^*$, and acts like $g' = g_{x_1} \dots g_{x_e}$ on it. In particular, the length of this word is at most k .

For all $x \in \Sigma \setminus \{x_1, \dots, x_d\}$ the restriction to the subtree $x\Sigma^*$ of g^f , where $f > 0$ is minimal such that g^f fixes x , has length at most $k - 1$, so is torsion by induction.

If g' has length k , then it is a product, in some order, of all the b'_i and a'_i with $\pi(b_i, a_i) = (b'_i, a'_i)$; since A is abelian, the order is unimportant. Sufficient applications of this step will lead us either to a situation where all a'_i fix the vertex d , and in that case g' has finite order at most $|B \times A|$; or to a situation where g' has length at most $k - 1$, in which case we may apply induction. $\square$

A subgroup G of $\text{Aut } \Sigma^*$ is *weakly branch* [Gri00] if it acts transitively on Σ^n for all $n \in \mathbb{N}$, and for all $\sigma \in \Sigma^*$ there exists $g \neq 1$ in G fixing pointwise $\Sigma^* \setminus \sigma\Sigma^*$. Weakly branch groups may not satisfy a non-trivial identity, so cannot be virtually nilpotent, so cannot have polynomial growth.

Note that there exist weakly branch groups of exponential growth — the simplest is probably $\langle a, b \rangle$ acting on $\{1, 2\}^*$, whose generators are defined by their decomposition $a = [1, b](1, 2)$ and $b = [1, a]$.

Branch groups are those weakly branch groups such that for every $n \in \mathbb{N}$ the subgroup generated by all elements acting only on some subtree $\sigma_1 \dots \sigma_n \Sigma^*$ has finite index in G . All known branch groups have intermediate growth.

Theorem 6.2. *Let G be a level-transitive regular mixer-splitter group of intermediate growth with A abelian. Then G is weakly branch.*

If furthermore G is torsion, then G is branch.

Proof. The subgroup G' is the kernel of the natural map $G \rightarrow A \times B$. Consider the subgroup $K = G''$; then $K \neq 1$ since then G would be solvable, and hence have polynomial or exponential growth.

Assume first $|X| \geq 3$. For any $k \in K$ we have $k' = [1, \dots, 1, k] \in K$; indeed for any $g_d \in G'$ we have some $[g_1, \dots, g_d] \in G'$, with at most one of $g_1, \dots, g_{d-1} \neq 1$. Taking commutators of two such expressions with other non-trivial element in different coordinates gives an arbitrary element $[1, \dots, 1, k]$ with $k \in K$.

If $|X| = 2$, then $|B| > |A|$ so we may follow similar considerations starting from elements in the kernel of ϕ_1^1 .

If G is torsion, then G'' has finite index in G . $\square$

7. FINAL REMARKS

In this final section I list a few remarks, extensions and open questions related to the growth of groups acting on trees.

1. It is worthwhile to stress where the inequalities are strict in the first part of Theorem 1.1's proof. First, the embedding $G_{k-1} \rightarrow G_k \wr_{\Sigma} A$ is not onto. Then in the decomposition process only some simplifications are performed. Finally the cancellations are assumed to occur independently — if they are correlated, then more cancellation occurs and the growth is smaller than predicted.
2. The extension of Theorem 6.2 to groups with non-abelian A should not pose any difficulty. However, is there a criterion to determine when such a group is branch? To see the difficulty, note that the groups 01 and 12 are branch, while 11 is only weakly branch. Among the examples with $d = 4$, it seems that all are branch except 012, 212, 103, 113, $1x1$.
3. Even though Theorem 1.1 covers all known constructions of groups of intermediate growth, it would be interesting to discover such groups of a completely different nature — for instance, simple groups, or finitely presented groups.
4. Andrew Brunner, Said Sidki and Ana Cristina Vieira [BSV99] studied extensively the subgroup G of $\text{Aut}\{1, 2\}^*$ generated by τ, μ defined by their decompositions $\tau = [1, \tau](1, 2)$ and $\mu = [1, \mu^{-1}](1, 2)$. They ask what the growth of G is, and this question is still open. Edmeia Da Silva [Sil01] showed that G does not contain any free subgroup; however, it is unknown whether G contains any free submonoids.
5. It should be possible to obtain quantitative estimates on the growth of G in Section 3, based on a study of the singularities of Γ on the unit circle imposed by (3).
6. We mentioned that the groups of polynomial growth are precisely the virtually nilpotent groups. Grigorchuk proved in [Gri89] that among residually- p groups, non-virtually-nilpotent groups have growth at least $e^{\sqrt{n}}$, and Alex Lubotzky and Avinoam Mann obtained the same result for residually-nilpotent groups—see [BG00] for more details.

If A is a p -group, the growth of G must therefore be at least $e^{\sqrt{n}}$. It seems reasonable to guess that this holds for all A , but a different argument may be needed.

It is not known whether there exist groups of growth strictly between polynomial and $e^{\sqrt{n}}$.

7. The analogous question for semigroups is well understood. Namely, given any two submultiplicative, superquadratic and subexponential functions γ, δ , there is a semigroup whose growth function is infinitely often larger than γ and infinitely often smaller than δ . George Bergman shows in [Ber78] that any polynomial growth rate $\gamma(n) = n^\alpha$ is possible with $\alpha \geq 2$, and Lev Shneerson shows in [Shn01] that the free semigroup with respect to a simple identity has intermediate growth of rate $e^{\sqrt{n}}$. See [Okn98] for semigroups of matrices with this property.
8. Groups of subexponential growth are amenable. It may be that all weakly branch groups are amenable, even those of exponential growth. There is no known example of a weakly branch group having Kazhdan's property (T).

9. The tree automorphisms corresponding to the $a \in A$ and $b \in B_k$ can be described by *transducer automata*. These are finite devices with states corresponding to the generators $A \cup B$, and transitions and activity corresponding to the decomposition map.

The nature of the automorphisms considered translates to the transitions being only from B states to A states, and from A states to the trivial state.

Said Sidki considered in [Sid00] a hierarchy of tree automorphisms, in which ours occupy the lowest layers. It might prove interesting to explore growth properties among higher groups in the Sidki ladder.

10. The proof of Section 3 extends *verbatim* to groups acting on irregular rooted trees $\Sigma_1 \times \Sigma_2 \times \dots$, with varying alphabets Σ_i and associated group A_i . In that setting G has subexponential growth as soon as $\sum_{n \in \mathbb{N}} \frac{1}{\#A_n}$ diverges. I would be hard pressed to find any usefulness to this generalization.

REFERENCES

- [Bar98] Laurent Bartholdi, *The growth of Grigorchuk's torsion group*, Internat. Math. Res. Notices **20** (1998), 1049–1054.
- [Bar01] Laurent Bartholdi, *Lower bounds on the growth of a group acting on the binary rooted tree*, Internat. J. Algebra Comput. **11** (2001), no. 1, 73–88.
- [Ben83] Michael Benson, *Growth series of finite extensions of $\mathbb{Z}^n$ are rational*, Invent. Math. **73** (1983), 251–269.
- [Ber78] George M. Bergman, *A note on growth functions of algebras and semigroups*, unpublished, 1978.
- [BG00] Laurent Bartholdi and Rostislav I. Grigorchuk, *Lie methods in growth of groups and groups of finite width*, Computational and Geometric Aspects of Modern Algebra (Michael Atkinson et al., ed.), London Math. Soc. Lect. Note Ser., vol. 275, Cambridge Univ. Press, Cambridge, 2000, pp. 1–27.
- [BG02] Laurent Bartholdi and Rostislav I. Grigorchuk, *On parabolic subgroups and Hecke algebras of some fractal groups*, to appear in Serdica Math. J., 2002, math.GR/9911206.
- [BŠ01] Laurent Bartholdi and Zoran Šunik, *On the word and period growth of some groups of tree automorphisms*, Comm. Algebra **29** (2001), no. 11, 4923–4964.
- [BSV99] Andrew M. Brunner, Said N. Sidki, and Ana Cristina Vieira, *A just nonsolvable torsion-free group defined on the binary tree*, J. Algebra **211** (1999), no. 1, 99–114.
- [Car21] Fritz Carlson, *über Potenzreihen mit ganzzahligen Koeffizienten*, Math. Z. **9** (1921), 1–13.
- [Fat06] Pierre Fatou, *Séries trigonométriques et séries de Taylor*, Acta Math. **30** (1906), 335–400.
- [FG85] Jacek Fabrykowski and Narain D. Gupta, *On groups with sub-exponential growth functions*, J. Indian Math. Soc. (N.S.) **49** (1985), no. 3-4, 249–256.
- [FG91] Jacek Fabrykowski and Narain D. Gupta, *On groups with sub-exponential growth functions. II*, J. Indian Math. Soc. (N.S.) **56** (1991), no. 1-4, 217–228.
- [GH90] Étienne Ghys and Pierre de la Harpe, *Sur les groupes hyperboliques d'après Mikhael Gromov*, Progress in Mathematics, vol. 83, Birkhäuser Boston Inc., Boston, MA, 1990, Papers from the Swiss Seminar on Hyperbolic Groups held in Bern, 1988.
- [Gri83] Rostislav I. Grigorchuk, *On the Milnor problem of group growth*, Dokl. Akad. Nauk SSSR **271** (1983), no. 1, 30–33.
- [Gri84] Rostislav I. Grigorchuk, *Degrees of growth of finitely generated groups and the theory of invariant means*, Izv. Akad. Nauk SSSR Ser. Mat. **48** (1984), no. 5, 939–985, English translation: Math. USSR-Izv. **25** (1985), no. 2, 259–300.
- [Gri89] Rostislav I. Grigorchuk, *On the Hilbert-Poincaré series of graded algebras that are associated with groups*, Mat. Sb. **180** (1989), no. 2, 207–225, 304, English translation: Math. USSR-Sb. **66** (1990), no. 1, 211–229.
- [Gri91] Rostislav I. Grigorchuk, *On growth in group theory*, Proceedings of the International Congress of Mathematicians, Vol. I, II (Kyoto, 1990) (Tokyo), Math. Soc. Japan, 1991, pp. 325–338.

- [Gri00] Rostislav I. Grigorchuk, *Just infinite branch groups*, New horizons in pro- p groups (Markus P. F. du Sautoy Dan Segal and Aner Shalev, eds.), Birkhäuser Boston, Boston, MA, 2000, pp. 121–179.
- [Gro81] Mikhael Gromov, *Groups of polynomial growth and expanding maps*, Inst. Hautes Études Sci. Publ. Math. (1981), no. 53, 53–73.
- [GS83a] Narain D. Gupta and Said N. Sidki, *On the Burnside problem for periodic groups*, Math. Z. **182** (1983), 385–388.
- [GS83b] Narain D. Gupta and Said N. Sidki, *Some infinite p -groups*, Algebra i Logika **22** (1983), no. 5, 584–589.
- [GS84] Narain D. Gupta and Said N. Sidki, *Extension of groups by tree automorphisms*, Contributions to group theory, Amer. Math. Soc., Providence, R.I., 1984, pp. 232–246.
- [Har00] Pierre de la Harpe, *Topics in geometric group theory*, University of Chicago Press, Chicago, IL, 2000.
- [Mil68a] John W. Milnor, *Growth of finitely generated solvable groups*, J. Differential Geom. **2** (1968), 447–449.
- [Mil68b] John W. Milnor, *Problem 5603*, Amer. Math. Monthly **75** (1968), 685–686.
- [Okn98] Jan Okniński, *Semigroups of matrices*, World Scientific Publishing Co. Inc., River Edge, NJ, 1998.
- [Par93] Walter Parry, *Growth series of Coxeter groups and Salem numbers*, J. Algebra **154** (1993), 406–415.
- [Sha89] Michael Shapiro, *A geometric approach to almost convexity and growth of some nilpotent groups*, Math. Ann. **285** (1989), 601–624.
- [Sha94] Michael Shapiro, *Growth of a $\mathrm{PSL}_2\mathbb{R}$ manifold group*, Math. Nachr. **167** (1994), 279–312.
- [Shn01] Lev M. Shneerson, *Relatively free semigroups of intermediate growth*, J. Algebra **235** (2001), no. 2, 484–546.
- [Sid00] Said N. Sidki, *Automorphisms of one-rooted trees: growth, circuit structure, and acyclicity*, J. Math. Sci. (New York) **100** (2000), no. 1, 1925–1943, Algebra, 12.
- [Sil01] Edméia Fernandes da Silva, *Uma família de grupos quase não-solúveis definida sobre árvores n -árias, $n \geq 2$* , Ph.D. thesis, Universidade de Brasília, 2001.
- [Sto96] Michael Stoll, *Rational and transcendental growth series for the higher Heisenberg groups*, Invent. Math. **126** (1996), no. 1, 85–109.
- [Šva55] Albert S. Švarc, *A volume invariant of coverings*, Dokl. Akad. Nauk SSSR (1955), no. 105, 32–34 (Russian).
- [Tit72] Jacques Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270.
- [Vov00] Taras Vovkivsky, *Infinite torsion groups arising as generalizations of the second Grigorchuk group*, Algebra (Moscow, 1998), de Gruyter, Berlin, 2000, pp. 357–377.
- [Wol68] Joseph A. Wolf, *Growth of finitely generated solvable groups and curvature of Riemannian manifolds*, J. Differential Geom. **2** (1968), 421–446.

E-mail address: laurent@math.berkeley.edu

DEPARTMENT OF MATHEMATICS, 970 EVANS HALL, CA94720-3840 BERKELEY, USA